

RI- SBIRT Screening Outcomes

September 2017-September 2019

Care Transformation Collaborative of R.I.

#1. Demographics by Site Setting

Figure 1a. Total # clients screened by site




Figure 1b. Gender by site




Figure 1c. Average age by site




Figure 1d. Hispanic ethnicity by site





Figure 1e. Race by site



#2. Screening Outcomes by Site Setting

Figure 2a. % of clients within each site who screened positive





Figure 2b. Percent of clients by site within each intervention modality



#3. Substance Use Outcomes

Figure 3a. Avg. # of reported alcohol use days in past month




Figure 3b. Avg. # of reported binge drinking days in past month




Figure 3c. Avg. # of reported marijuana use days in the past month




Figure 3d. Percent of clients who reported opioid use past month at intake and 6 month follow-up

Intake





Figure 3e. Avg. # of reported illegal drug use days in the past month



Discussion

- 1. What reflections/surprises do you have to share about the data presented?
- 2. What has the experience been like for you and your site to have SBIRT screenings available?
- 3. Moving into year 3 of SBIRT, are there any opportunities for new site types or changes to where we are currently screening?
- 4. How can we make SBIRT sustainable with SAMHSA grant resources?



CELEBRATING 1 YEAR OF RI-SBIRT!

September 2017 - September 2018

7,858 Screens Completed

1,460 interventions



9 out of 10 people at risk received an intervention.

Clients who received an intervention for drug use reported a significant decrease in the average number of illegal drug use days at follow-up.



■ Intake ■ 6 month follow-up

5.9

Avg. # days in past month used illegal drugs

Clients who received an intervention for risky alcohol use reported a <u>significant decrease</u> in the average number of alcohol use days at follow-up.



■ Intake ■ 6 month follow-up

16.9 8.1

Avg. # days in past month used alcohol

1 of every 2 (53%) clients who reported illegal drug use at intake reported no use 6 months later.




Of those selected for follow-up, only 15% of clients who reported opiate use at intake reported use at follow-up.

20 clients at intake



3 clients at follow-up


Clients who reported marijuana use at intake reported significantly fewer marijuana use days at follow-up.



Avg. # days in past month used marijuana

There was a <u>significant decrease</u> (49%) in the number of clients who reported past month alcohol use from intake to follow-up.

2 out of 5 (40%) clients who reported binge drinking in the month before intake reported no binge drinking at follow-up.



Murphy, L., Bassett, S.S., & Stein, L.A.R. (2018). Celebrating one year of RI-SBIRT [Handout]. Kingston, RI: University of Rhode Island.

RT = referral to treatment; BT = brief treatment; BI = brief intervention